

Physics 234: Practice Lab Test

1. How many of the numbers $1, 2, \dots, 1000$ are perfect squares?

31

How many are perfect cubes?

10

(Hint: you can solve this question without using either `sqrt` or `pow`.)

2. Consider an unbounded square grid of points spaced by $\Delta x = 0.1$ and $\Delta y = 0.1$. How many points lie inside the diamond $|x| + |y| = 2$?

761

How many points lie inside the circle $x^2 + y^2 = 4$?

1245

(Hints: (i) don't count the points on the boundary; (ii) you may be better off reformulating the problem so that you can use integer rather than floating-point types in your code.)

3. Here's a variant of *Buffon's needle problem*. Imagine placing a randomly-directed straight line segment of unit length in the plane so that one end lies on the x -axis somewhere between -1 and 1 . What is the probability P that the line segment intersects the y -axis?

Evaluate the probability as follows:

- Set integers counter N and C to zero.
- Choose three random numbers $\xi, \xi', \xi'' \in [-1, 1]$.
- If the point (ξ', ξ'') lies in the annulus defined by $0.001 < (\xi')^2 + (\xi'')^2 \leq 1$ then construct a unit vector

$$\hat{\mathbf{n}} = \frac{(\xi', \xi'')}{\sqrt{(\xi')^2 + (\xi'')^2}},$$

position the end-points at $\mathbf{r}_1 = (\xi, 0)$ and $\mathbf{r}_2 = \mathbf{r}_1 + \hat{\mathbf{n}}$, and increment N by one.

- If the line segment connecting \mathbf{r}_1 and \mathbf{r}_2 crosses the y -axis then increment C by one.
- Repeat from (b) until N numbers in the tens of millions.
- Report the result $P \doteq C/N$.

0.318

4. Consider the sequence $(a_n) = (a_0, a_1, a_2, \dots)$ whose elements are the partial sums

$$a_n = \sum_{k=0}^n (z^k + 2z^{k+1} + 3z^{2k}).$$

This sequence converges linearly and attains the limit

$$L = \lim_{n \rightarrow \infty} a_n = \frac{1}{1-z} + \frac{2z}{1-z} + \frac{3}{1-z^2} = \frac{4+3z+2z^2}{1-z^2}.$$

In the case of $z = 0.1$, the first three terms are $a_0 = 4.2$, $a_1 = 4.35$, and $a_2 = 4.3623$. What is the smallest value p for which a_p agrees with L to within 1 part in 10^6 .

$$p = \boxed{5} \quad a_p = \boxed{4.3636350303}$$

Generate a new sequence (A_n) from (a_n) using the Aitken transformation:

$$A_n = a_{n+1} + \left(\frac{1}{a_{n+2} - a_{n+1}} - \frac{1}{a_{n+1} - a_n} \right)^{-1}$$

What is the smallest value q for which A_q agrees with L to within 1 part in 10^6 .

$$q = \boxed{1} \quad A_q = \boxed{4.36363341443634}$$